

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 86118045.3

(22) Date of filing: 24.12.86

(51) Int. Cl.<sup>3</sup>: C 07 D 211/26

C 07 D 401/12, C 07 D 401/0- 6  
 C 07 D 211/22, C 07 D 211/1- 4  
 C 07 D 211/32, C 07 D 211/7- 0  
 C 07 D 211/94, C 07 D 401/0- 4  
 C 07 D 401/14, C 07 D 405/1- 2

(30) Priority: 27.12.85 JP 293885/85

(43) Date of publication of application:  
22.07.87 Bulletin 87/30

(84) Designated Contracting States:  
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: Eisai Co., Ltd.  
6-10, Kolshikawa 4-chome Bunkyo-ku  
Tokyo 112(JP)

(72) Inventor: Sugimoto, Hachiro  
3073-13, Kashiwadamachi  
Ushiku-shi Ibaraki(JP)

(72) Inventor: Nakamura, Takaharu  
4-9, Tsukushino 2-chome  
Abiko-shi Chiba(JP)

(72) Inventor: Tsuchiya, Yutaka  
35-16, Sakaecho 2-chome  
Ushiku-shi Ibaraki(JP)

(72) Inventor: Sugumi, Hiroyuki  
Shizanryo 19-13, Kasuga 4-chome Yatabemachi  
Tsukuba-gun Ibaraki(JP)

(72) Inventor: Higurashi, Kunizou  
4-3, Midori 2-chome  
Sumida-ku Tokyo(JP)

(72) Inventor: Karibe, Norio  
Shizanryo 19-13, Kasuga 4-chome Yatabemachi  
Tsukuba-gun Ibaraki(JP)

(72) Inventor: Yamanishi, Yoshiharu  
2079, Wakashibacho  
Ryugasaki-shi Ibaraki(JP)

(72) Inventor: Ogura, Hiroo  
1125-5, Ohaza Nagakuni  
Tsuchiura-shi Ibaraki(JP)

(72) Inventor: Araki, Shin  
Kashiwa Mansion 401 11-6 Takezono 2-chome  
Sakuramura Niihari-gun Ibaraki(JP)

(72) Inventor: Kubota, Atsuhiko  
New life Namiki 406 15-1, Namiki 4-chome  
Sakuramura Niihari-gun Ibaraki(JP)


(72) Inventor: Ohtake, Michiko  
3369, Suwamachi  
Mitsukaido-shi Ibaraki(JP)

(72) Inventor: Yamatsu, Kiyomi  
23-7, Imaizumidai 7-chome  
Kamakura-shi Kanagawa(JP)

(74) Representative: Hansen, Bernd, Dr.rer.nat. et al,  
Hoffmann, Eitle & Partner Patentanwälte Arabellastrasse  
4  
D-8000 München 81(DE)

(54) Piperidine derivative, its use and pharmaceutical composition containing it.

(57) A novel piperidine derivative as defined by the formula cerebrovascular disease is disclosed.  
(I), including a salt thereof,




wherein R<sup>1</sup>, X, A and R<sup>2</sup> have the aforementioned meanings.  
Further, pharmaceutical compositions containing the same  
and the use of the piperidine derivatives for the making of  
such compositions preventing dementias and sequelae of

Piperidine Derivative, its use and Pharmaceutical  
Composition containing it

---

The present invention relates to piperidine derivatives having excellent actions as medicines. The invention relates more particularly to piperidine derivatives of the following general formula (I) or pharmacologically allowable salts thereof:



wherein  $R^1$  denotes a univalent group derived from one selected among substituted or unsubstituted benzene, pyridine, pyrazine, indole, anthraquinone, quinoline, substituted or unsubstituted phthalimide, homophthalimide, pyridinecarboxylic acid imide, pyridine N-oxide, pyrazinedicarboxylic acid imide, naphthalenedicarboxylic acid imide, substituted or unsubstituted quinazolinedione, 1,8-naphthalimide, bicyclo [2.2.2] octo-5-ene-2,3-dicarboxylic acid imide and pyromerylimide,

X denotes a group of the formula  $-(CH_2)_n-$ , a group of the formula  $-O(CH_2)_n-$ , a group of the formula  $-S(CH_2)_n-$ , a group of the formula  $-NH(CH_2)_n-$ , a group of the formula  $-SO_2NH(CH_2)_n-$ , a group of the formula  $-NH-\overset{\overset{O}{||}}{C}-(CH_2)_n-$ , a group of the formula  $-NH(CH_2)_n-\overset{\overset{O}{||}}{C}-$ , a group of the formula


$\begin{array}{c} \text{O} \\ || \\ -\text{C}-\text{O}(\text{CH}_2)_n- \end{array}$ , a group of the formula  $-\text{CH}_2\text{NH}(\text{CH}_2)_n-$ , a group


of the formula  $\begin{array}{c} || \quad | \\ -\text{C}-\text{N}-(\text{CH}_2)_n- \\ \text{O} \quad \text{R}^3 \end{array}$  (in all the above formulas,

5  $n$  is an integer of 1 through 7 and  $\text{R}^3$  represents a lower alkyl group or a benzyl group), a group of the formula



$\begin{array}{c} \text{CH}_3 \\ | \\ -\text{O}-\text{CH}_2\text{CH}_2\text{CH}- \end{array}$ , a group of the formula  $\begin{array}{c} \text{CH}_3 \\ | \\ -\text{O}-\text{CHCH}_2\text{CH}_2- \end{array}$ , a group of the formula  $-\text{O}-\text{CH}_2\text{CH}_2\text{CH}=$  or a group of the formula

10  $\begin{array}{c} \text{OH} \\ | \\ -\text{O}-\text{CH}_2-\text{CH}-\text{CH}_2- \end{array}$ ,

the ring A denotes a group of the formula ,

a group of the formula , a group of the formula

15

 or a group of the formula , and

$\text{R}^2$  denotes a hydrogen atom, a lower alkyl group, a substituted or unsubstituted benzyl group, a substituted or unsubstituted benzoyl group, a pyridyl group, a 2-hydroxyethyl group, a pyridylmethyl group or a group of the formula  $\begin{array}{c} \text{OH} \\ | \\ \text{Z}-\text{C}_6\text{H}_4-\text{CH}- \end{array}$  (wherein  $\text{Z}$  represents a halogen atom).


The lower alkyl group in the definition of the formula  
25 (I) means a straight-chain or branched alkyl group having

1 to 6 carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, 1-methylpropyl, tert-butyl, n-pentyl, 1-ethylpropyl, isoamyl or n-hexyl. Preferable lower alkyl groups include a methyl group, ethyl group, etc.

- 5 The lower alkoxy group means a group derived from the above lower alkyl group.

The univalent group derived from substituted or unsubstituted benzene in the definition of  $R^1$  means specifically the following:


- 10 (1) Univalent group of the formula



- wherein n is an integer of 1 through 3 and D denotes a phenyl group optionally substituted by a group or two or three same or different groups selected among a hydrogen atom, a lower alkyl group, a nitro group, a lower alkoxy group, an alkylenedioxy group formed between adjacent carbon atoms in arbitrary positions, a cyano group, a halogen atom, an amino group, a monoalkylamino or dialkylamino group, a lower alkoxycarbonyl group, a trifluoromethyl group, a formyl group, a hydroxy group (hydroxyl group), a lower alkylthio group, a lower alkylsulfinyl group, a lower alkylsulfonyl group, a lower alkylsulfoxide group, a lower alkylcarbonyl group, a methoxymethylthio group, a halogenomethylthio group, a cycloalkylsulfonyl group, a phenyl group, a cycloalkylthio group and a cyclohexenyloxy
- 15  
20  
25

group.

(2) Univalent group of the formula




5 wherein G denotes a group of the formula  $\text{-}\overset{\text{O}}{\parallel}\text{C-}$ , a group of the formula  $\text{-O-}\overset{\text{O}}{\parallel}\text{C-}$ , a group of the formula  $\text{-O-}$ , a group of the formula  $\text{-CH}_2\text{-NH-}\overset{\text{O}}{\parallel}\text{C-}$ , a group of the formula  $\text{-CH}_2\text{-O-}$ ,  
 10 a group of the formula  $\text{-CH}_2\text{-SO}_2\text{-}$ , a group of the formula  $\text{-CH-}$  or a group of the formula  $\text{-CH}_2\text{-}\overset{\text{O}}{\underset{\uparrow}{\text{S}}}\text{-}$ , E denotes a carbon  
 $\text{OH}$   
 atom or a nitrogen atom, and D has the same meaning as  
 15 defined in (I).

In the univalent group derived from substituted or unsubstituted phthalimide in the definition of  $R^1$ , the examples of the preferable substituent include a nitro group, an amino group, a halogen group, a lower alkyl group, a  
 20 lower alkoxy group, a hydroxy group, a benzoyl group, a phenylcarbonyl group, a phenylcarbonylamino group, a lower alkylcarbonylamino group, a hydroxycarbonyl group, a benzylaminocarbonyl group and a dialkylaminocarbonyl group. The univalent group may be substituted by two or more same  
 25 or different substituents if necessary.

In the univalent group derived from substituted or unsubstituted quinazolinedione in the definition of R<sup>1</sup>, the examples of the preferable substituent include a lower alkyl group and a halogen group.

5 Hereinafter preferred methods of making the present compounds are disclosed which methods are fully encompassed by the present invention:

Preparation Method A





wherein Hal denotes a halogen atom, and R<sup>1</sup>, X, R<sup>2</sup> and the ring A have the same meaning as defined above.

Namely, a compound of the general formula (II) (wherein  
 15 Hal denotes a chlorine atom, bromine atom, iodine atom, etc. and among them the bromine atom is most preferable) and a piperidine derivative of the general formula (III) are subjected to a condensation reaction by a conventional method, preferably in the presence of a base such as sodium  
 20 hydrogencarbonate, sodium carbonate, potassium carbonate or triethylamine, to obtain the final compound (I). In this case, as an organic solvent is used, for example, benzene, toluene, ethanol, butanol or dimethylformamide (DMF).

Preparation Method B

25 [In the case where X denotes a group of the formula






Namely, an acid halogenide of the general formula (IV) is allowed to react with a piperidine derivative of the general formula (V) in an organic solvent such as chloroform, benzene, toluene, dioxane, tetrahydrofuran or dimethylformamide (DMF), in the presence of a desalting agent such as sodium carbonate, potassium carbonate, sodium hydroxide, potassium hydroxide or triethylamine, with ice-cooling, at room temperature or with heating, to easily obtain the compound (VI), one of the final compounds.

#### Preparation Method C


[In the case where X denotes a group of the formula  $-\text{SO}_2-\text{NH}(\text{CH}_2)_n-$  in the general formula (I)]



Procedure similar to that of Preparation Method B is used to obtain the compound (IX), one of the final compounds.


Preparation Method D

[In the case where  $R^1$  denotes a univalent group derived  
 5 from imide selected among substituted phthalimide,  
 homophthalimide, pyridinecarboxylic acid imide,  
 pyrazinedicarboxylic acid imide, naphthalenedicarboxylic  
 acid imide, 1,8-naphthalimide, bicyclo [2.2.2] octo-5-ene-  
 2,3-dicarboxylic acid imide and pyromerylimide and X  
 10 denotes a lower alkylene group in the general formula (I)]



wherein  $n$  is an integer of 1 through 7 and B denotes a residue after the removal of a group of the formula






in all the above-defined R<sup>1</sup>.

5       Namely, an acid anhydride of the general formula (X) and a piperidine derivative of the general formula (XI) are subjected to a condensation reaction by a conventional method to obtain the compound (XII), one of the object substances.

10       The reaction is carried out with application of heat in an organic solvent such as, for example, ethanol, butanol, dioxane, dimethylformamide (DMF) or acetic anhydride.

#### Preparation Method E

15       [In the case where R<sup>1</sup> denotes a univalent group derived from substituted quinazolinedione and X denotes a lower alkylene group]




15  $\text{R}^5$  and  $\text{R}^6$  are hydrogen, or a substituent such as a lower alkyl and a halogen.

Namely, a diester of the general formula (XIII) is allowed to react with a piperidine derivative of the general formula (XIV), with application of heat, in a suitable solvent which does not participate in the reaction or in the absence of the solvent, to obtain the quinazolone compound (XV), one of the object substances.

#### Preparation Method F

25 [In the case where  $\text{R}^1$  denotes a univalent group derived from substituted or unsubstituted benzene and X denotes a group of the formula  $-\text{O}(\text{CH}_2)_n-$  in the general formula (I)]





5  
 10  $\text{R}^1$  is hydrogen or a phenyl defined above.  $p$  is zero or an integer of 1 to 3.

Namely, a phenol derivative of the general formula (XVI) and a halogen compound of the general formula (XVII) are subjected to a condensation reaction by a conventional  
 15 method to obtain the compound (XVIII), one of the final compounds.

The reaction is carried out in a solvent such as, for example, tetrahydrofuran or dimethylformamide (DMF), in the presence of NaH or NaOH, at room temperature or with heating,  
 20 thereby obtaining a good result.


#### Preparation Method G

[In the case where  $\text{R}^1$  denotes a univalent group derived from substituted or unsubstituted benzene and X denotes a group of the formula  $\text{---S(CH}_2\text{)}_n\text{---}$  in the general formula (I)]



Procedure similar to that of Preparation Method F is used to obtain the compound (XXI), one of the object substances.

#### Preparation Method H



wherein  $R^1$ ,  $X$ ,  $R^2$  and the ring  $A$  have the same meaning as defined above.

Namely, a compound of the general formula (XXII) and a piperidine derivative of the general formula (III) are subjected to a condensation reaction preferably using a desalting agent such as triethylamine, N-methylmorpholine or N,N'-dimethylaniline to obtain the object substance (I).

In this case, benzene, toluene, tetrahydrofuran, dimethylformamide or dioxane is used as a solvent.

Various attempts have been made to treat for middle-  
ager dementia, senile dementia and so on with medicines. At  
5 present, however, there is no medicine which is considered  
to be drastically effective for the diseases. Considered  
to be effective at present are anticholinesterase agents  
(example; physostygmine). The physostygmine, however,  
suffers disadvantages: action of short duration, strong  
10 side effects, and so forth.

Accordingly, the inventors have been making many  
intensive studies over a long term of years in order to  
develop medicines having actions of long duration and being  
high in safety.

15 As a result, they have discovered that the piperidine  
derivatives of the general formula (I) can attain the  
desired end.

Specifically, the compounds of the structural formula  
(I) according to the invention have the following important  
20 features. They have strong and highly-selective  
antiacetylcholinesterase activities. Furthermore, they  
increase the amount of acetylcholine in the brain and are  
effective for the model of retentive disorder. In addition,  
they have actions of long duration and are high in safety,  
25 compared with the physostygmine heretofore in use in the

field. Thus, the invention is of great value.

Namely, the compounds of the invention are piperidine derivatives of the following general formula (I) or pharmacologically allowable salts thereof:



wherein R<sup>1</sup> denotes a univalent group derived from one selected among substituted or unsubstituted benzene, pyridine, pyrazine, indole, anthraquinone, quinoline, substituted  
 10 or unsubstituted phthalimide, homophthalimide, pyridinecarboxylic acid imide, pyridine N-oxide, pyrazinedicarboxylic acid imide, naphthalenedicarboxylic acid imide, substituted or unsubstituted quinazolidinedione, 1,8-naphthalimide, bicyclo [2.2.2] octo-5-ene-2,3-dicarboxylic acid imide and  
 15 pyromerylimide,

X denotes a group of the formula  $-(CH_2)_n-$ , a group of the formula  $-O(CH_2)_n-$ , a group of the formula  $-S(CH_2)_n-$ , a group of the formula  $-NH(CH_2)_n-$ , a group of the formula  $-SO_2NH(CH_2)_n-$ , a group of the formula  $-NH-\overset{\overset{O}{\parallel}}{C}-(CH_2)_n-$ , a  
 20

group of the formula  $-NH(CH_2)_n-\overset{\overset{O}{\parallel}}{C}-$ , a group of the formula

$-\overset{\overset{O}{\parallel}}{C}-O(CH_2)_n-$ , a group of the formula  $-CH_2NH(CH_2)_n-$ , a group

of the formula  $\begin{array}{c} \text{-C-N-(CH}_2\text{)}_n\text{-} \\ \parallel \quad | \\ \text{O} \quad \text{R}^3 \end{array}$  (in all the above formulas,


n is an integer of 1 through 7 and R<sup>3</sup> represents a lower alkyl group or a benzyl group), a group of the formula


5

$\begin{array}{c} \text{CH}_3 \\ | \\ \text{-O-CH}_2\text{CH}_2\text{CH-} \end{array}$ , a group of the formula  $\begin{array}{c} \text{CH}_3 \\ | \\ \text{-O-CHCH}_2\text{CH}_2\text{-} \end{array}$ , a group of the formula  $\text{-O-CH}_2\text{CH}_2\text{CH=}$  or a group of the formula



$\begin{array}{c} \text{OH} \\ | \\ \text{-O-CH}_2\text{-CH-CH}_2\text{-} \end{array}$ ,

10

the ring A denotes a group of the formula ,

a group of the formula , a group of the formula

15

 or a group of the formula , and

20

R<sup>2</sup> denotes a hydrogen atom, a lower alkyl group, a substituted or unsubstituted benzyl group, a substituted or unsubstituted benzoyl group, a pyridyl group, a 2-hydroxyethyl group, a pyridylmethyl group or a group of the formula  $\begin{array}{c} \text{OH} \\ | \\ \text{Z} \text{---} \text{C}_6\text{H}_4 \text{---} \text{CH-} \end{array}$  (wherein Z represents a halogen atom).

25

Therefore purposes of the invention are to provide a novel compound which is effective to various types of dementias and sequelae of cerebrovascular diseases, then methods for preparation of the compound and a pharmaceutical composition comprising the compound as the effective ingredient.

The pharmacologically acceptable salt to use in the invention includes for instance an inorganic salt with hydrochloride, sulfate, hydrobromate or phosphate and an organic salt with formate, acetate, trifluoroacetate, maleate, tartrate, methanesulfonate, benzenesulfonate or toluenesulfonate.

The compound of the invention is effective to treatment, prevention, remission and improvement of various types of senile dementia, especially senile dementia of the Alzheimer type or the Alzheimer's disease, the disturbance of attention, aphasia, hypobulia, the emotional disorder, the memory disorder, the hallucinatory-paranoid state and the abnormal behavior, accompanying and following cerebrovascular diseases such as cerebral apoplexy (cerebral hemorrhage, cerebral infarction), sequelae of encephalitis and cerebral palsy.

Further, the compound of the invention has a strong, highly selective anticholinesterase activity and is eventually useful as a medicine based on the activity.

When the compound is used as the medicine, it may be orally or parenterally administered. It is parenterally administered in the form of an intravenous, hypodermic or intramuscular injection or a suppository. It may be administered also in the form of a sublingual tablet. A dose of the administration depends on conditions of a patient, such as age, sex, a body weight and sensitivity, a method of the administration such as times and intervals, properties, preparation and kinds



of the medicine, kinds of effective ingredients, and in case another treatment is also effected simultaneously, the kind, frequency and intended effects of the treatment.

In general, a dose of the administration is about 0.1  
5 to 300 mg, preferably about 1 to 100 mg, per an adult a day. The administration with the amount is made one to four times a day.

When the compounds of the invention are prepared into medicines, they are prepared into medicines in the form of  
10 injections, suppositories, sublingual tablet, tablets, capsules, etc. using ordinary carrier by a conventional method in the technical field of preparation.

In the preparation of injections, pH regulator, buffer, suspending agent, dissolution adjuvant, stabilizer,  
15 preservative, etc. are added to the principal ingredient when required, and intravenous, hypodermic and intramuscular injections are prepared by a conventional method. In this case, the injections may be frozen and dried, if necessary, by a conventional method.

20 Examples of the suspending agent include methyl cellulose, polysorbate 80, hydroxyethyl cellulose, gum arabic, tragacanth powder, carboxymethyl cellulose sodium, polyoxyethylene sorbitane monolaurate, etc.

Examples of the dissolution adjuvant include  
25 polyoxyethylene-hardened castor oil, polysorbate 80, amide nicotinate, polyoxyethylene sorbitane monolaurate, castor

oil fatty acid ethyl ester, etc.


Examples of the stabilizer include sodium sulfite, sodium metasilicate, ether, etc. Examples of the preservative include methyl paraoxybenzoate, ethyl paraoxybenzoate,  
 5 sorbic acid, phenol, cresol, chlorocresol, etc.

The representative compounds of the invention will be described hereinafter by way of examples. It goes without saying, however, that the representative compounds are for purpose of help to the understanding of the invention and are  
 10 not intended as a definition of the limits of the invention.

The values of NMR in the following examples are those in free substances.

Example 1

1-benzyl-4-[γ-(4-nitrophenoxy) propyl] piperidine . hydrochloride



0.8 g of p-nitrophenol and 1.4 g of N-benzyl-4-(γ-bromopropyl)-piperidine are dissolved in 20 ml of dimethylformamide (DMF). To the mixed solution is added,  
 20 little by little, with stirring at room temperature, 0.3 g of 60% sodium hydride.

Hereafter, the resulting mixture is stirred for 2 hours at room temperature and is further stirred for about 6 hours and 30 minutes at 70 to 80 °C. After the solvent is  
 25 distilled off under reduced pressure, an aqueous chloroform-

5% caustic soda solution is added to the residue followed by shaking sufficiently with a separating funnel to separate out a chloroform layer.

The chloroform layer is washed with a saturated saline solution, which is then dried over magnesium sulfate. Chloroform is distilled off under reduced pressure, and the resulting residue is purified using a silica gel column. The distillation is performed using 2% methanol-chloroform. After distilling off the solvent under reduced pressure, the residue is dissolved in ethyl acetate followed by adding a 10% hydrochloric acid-ethyl acetate solution to separate out crystals. Upon recrystallization from ethanol-water-ethyl ether, 1.9 g (yield: 84.1%) of the titled compound having the following physical properties is obtained.


Melting point ( $^{\circ}\text{C}$ ): 229-230

Elemental analytical values:  $\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_3 \cdot \text{HCl}$

	C	H	N
Theoretical value (%)	64.52	6.96	7.17
Found value (%)	64.21	7.03	7.06

## Example 2

N-[4'-(1'-benzylpiperidine) ethyl]-4-benzylsulfonylbenzamide . hydrochloride



3 g of 1-benzyl-4-aminoethylpiperidine and 2.8 g of

triethylamine are added to 100 ml of anhydrous tetrahydrofuran. 4.1 g of 4-benzylsulfonylbenzoylchloride is mixed with 50 ml of anhydrous tetrahydrofuran with ice-cooling and stirring, which is then added dropwise to the above mixed solution for about 20 minutes. The resulting mixture is stirred for about 20 minutes at room temperature and is further refluxed for about 20 minutes. Then, tetrahydrofuran is distilled off under reduced pressure. The resulting residue is purified using a column in a similar manner as in Example 1 and is formed into a hydrochloride by a conventional method to obtain 3.55 g of the titled compound (yield: 49.4%).


Melting point (°C): 187-188

Elemental analytical values:  $C_{28}H_{32}N_2O_3S \cdot HCl$

	C	H	N
Theoretical value (%)	65.55	6.48	5.46
Found value (%)	64.64	6.27	5.36
3/10 (H <sub>2</sub> O)	64.86	6.53	5.46

### Example 3

N-[4'-(1'-benzylpiperidine) ethyl]-4-isonicotinic acid amide . hydrochloride



(Preparation Example 1)

2.4 g of isonicotinic acid chloride · hydrochloride is

added little by little to 4.4 g of 1-benzyl-4-aminoethylpiperidine, 5.6 g of potassium carbonate and 50 ml of dioxane while they are cooled with ice and stirred. After the reaction for about 1 hour at room temperature, 5 dioxane is distilled off under reduced pressure. To the resulting residue are added 50 ml of water and 20 ml of a 5% aqueous NaOH solution to alkalify the residue. The alkaline residue is then extracted with chloroform. After washing with water, the resulting chloroform layer is dried 10 over potassium carbonate, and chloroform is distilled off under reduced pressure. The resulting residue is purified with 2% methanol-chloroform-based and 5% methanol-chloroform-based solvents by the use of a silica gel column, and is formed into a hydrochloride using a 10% hydrochloric 15 acid-ethyl acetate solution to obtain 5.1 g of the titled compound (yield: 70.0%).

(Preparation Example 2)

4.7 g of isonicotinic acid is dissolved in a mixed solution of dimethylformamide-tetrahydrofuran (1/1). To the 20 mixture are added, at -30 to -15 °C, 3.85 g of N-methylformalin and 4.13 g of ethyl chlorcarbonate. The resulting mixture is stirred for 5 minutes at -15 °C. Dimethylformamide-tetrahydrofuran solution of 8.33 g of N-benzylpiperidylethylamine is added to the reaction 25 solution followed by stirring for 2 hours at 0 °C and then


1 hour at room temperature. After removing a precipitate by filtration, the resulting filtrate is concentrated, and the residue is dissolved in chloroform. After washing with caustic soda and water, the chloroform layer is dried over magnesium sulfate, and chloroform is distilled off. The resulting oily residue is purified using a silica gel chromatograph to obtain 9.5 g of the titled compound (yield: 77.0%).

NMR( $\delta$  value, DMSO):

1.14-2.04 (9H, m), 2.8-2.9 (2H, bd),  
 3.36-3.56 (2H, m), 3.44 (2H, s),  
 6.2 (1H, bt), 7.24 (5H, s), 7.54 (2H, m),  
 8.68 (2H, m)

#### Example 4

N-methyl-N-[4'-(1'-benzylpiperidine) ethyl]-4-benzylsulfonylbenzamide . hydrochloride



5.9 g of 4-benzylsulfonylbenzoylchloride is added little by little to 4.6 g of 1-benzyl-4-(N'-methylaminoethyl) piperidine, 5 g of triethylamine and 40 ml of chloroform while they are cooled with ice and stirred. After the reaction for 12 hours at room temperature, 20 ml of water and 20 ml of a 5% aqueous NaOH solution are added to the reaction solution followed by shaking sufficiently with a

separating funnel to separate out a chloroform layer. After washing with water, the chloroform layer is dried over magnesium sulfate, and chloroform is distilled off under reduced pressure. The resulting residue is purified using a column in a similar manner as in Example 1 and is formed into a hydrochloride. Upon recrystallization from ethanol, 8.2 g of the titled compound is obtained (yield: 78.1%).


Melting point (°C): 200-201

Elemental analytical values: C<sub>29</sub>H<sub>34</sub>N<sub>2</sub>O<sub>3</sub>S

	C	H	N
Theoretical value (%)	66.08	6.69	5.31
Found value (%)	66.12	6.67	5.21

#### Example 5

N-methyl-N-[4'-(1'-benzylpiperidyl) ethyl] isonicotinic acid amide hydrochloride



3.48 g of N-benzyl-4-(N'-methyldaminoethyl) piperidine and 4.6 g of potassium carbonate are added to a mixed solution of 40 ml of chloroform and 10 ml of water. To the mixture is added, little by little, with ice-cooling and stirring, 3.2 g of isonicotinic acid chloride · hydrochloride. After stirring for 1 hour at room temperature, 20 ml of

- 23 -

water and 10 ml of an aqueous 1N-NaOH solution are added to the reaction solution, and a chloroform layer is separated out. After washing with water, the chloroform layer is dried over magnesium sulfate.

- 5 Chloroform is distilled off under reduced pressure to obtain 4.3 g of an oily matter. The oily matter is purified using a silica gel column in a similar manner as in Example 1 and is formed into a hydrochloride.


Upon recrystallization from acetone-ethanol, 4.0 g of  
10 the titled compound is obtained (yield: 72.0%).

Elemental analytical values:  $C_{21}H_{27}N_3O \cdot 2HCl \cdot 1/2H_2O$

	C	H	N
Theoretical value (%)	60.14	7.21	10.12
Found value (%)	60.02	7.01	10.16

15 Example 6

N-methyl-N-[4'-(1'-benzylpiperidine) ethyl]-4-cyclopentylsulfonylbenzamide . hydrochloride



- 1.1 g of N-benzyl-4-(N'-methylaminoethyl) piperidine and 1.4 g of potassium carbonate are added to a mixed solution of 20 ml of chloroform and 5 ml of water. While stirring the mixture at room temperature, a solution in  
25 which 1.16 g of 4-cyclopentylsulfonylchloride is dissolved



in 20 ml of chloroform is added dropwise to the mixture. After stirring for 2 hours at room temperature, 10 ml of water and 5 ml of an aqueous 1N-NaOH solution are added to the reaction solution, and a chloroform layer is separated  
 5 out. After washing with water, the chloroform layer is dried over magnesium sulfate.

Chloroform is distilled off under reduced pressure to obtain a crude product. The crude product is purified using a silica gel column in a similar manner as in Example 1 and  
 10 is formed into a hydrochloride.

Upon recrystallization from ethanol-ether, 1.9 g of the titled compound is obtained (yield: 80.0%).

Melting point (°C): 234-236 (decomposition)


Elemental analytical values:  $C_{27}H_{36}N_2O_3S \cdot HCl \cdot 1/2H_2O$

15		C	H	N
	Theoretical value (%)	63.08	7.45	5.45
	Found value (%)	63.10	7.25	5.40

#### Example 7

N-[2-(N'-benzylpiperidino-4) ethyl]-4-nitrophthalimide

20 hydrochloride



1.0 g of 4-nitro phthalic anhydride and 1.1 g of 4-(2-aminoethyl)-benzylpiperidine are added to 30 ml of dioxane. The resulting mixture is heated and refluxed for 2 hours.

5 The reaction solution is added with 50 ml of water followed by extracting with chloroform. The resulting chloroform layer is dried over magnesium sulfate, which is then concentrated to dryness under reduced pressure. The resulting residue is purified with a 5% ethanol-chloroform-  
10 based solvent by the use of a silica gel column to obtain 1.3 g of the object substance. The object substance is formed into a hydrochloride using a 10% hydrochloric acid-ethyl acetate solution. Upon recrystallization from acetone-isopropyl ether, 0.98 g of the titled compound is obtained  
15 (yield: 45.1%).


Melting point (°C): 224-227

Elemental analytical values:  $C_{22}H_{23}N_3O_4 \cdot HCl$

	C	H	N
Theoretical value (%)	61.47	5.63	9.77
20 Found value (%)	61.35	5.69	10.01

#### Example 8

N-[2-(N'-benzylpiperidino-4) ethyl]-2,3-pyrazinedicarboxylic acid imide




7.5 g of 2,3-pyrazinedicarboxylic anhydride and 12.0 g of 4-(2-aminoethyl)-benzylpiperidine are stirred for 10 minutes at 120 °C to form a brown tar-like mixture. After allowing to stand for cooling, 7 ml of acetic anhydride is added dropwise to the brown tar-like mixture at 80 °C followed by stirring for about 30 minutes at 100 °C. The resulting tar-like substance is added with and dissolved in 20 ml of chloroform and is purified using a silica gel column in a similar manner as in Example 1. Since the object substance is not well separated out, the above tar-like substance is further purified by distillation with benzene and a 10% ethanol-benzene-based solvent, to obtain 5.9 g of the object compound as a yellow-brown oily matter (yield: 33.6%).

NMR ( $\delta$  value, DMSO):

1.27-2.02 (9H, m), 2.8-2.9 (2H, bd),  
3.46 (2H, s), 3.82 (2H, t), 7.26 (5H, s),  
8.87 (2H, s)

#### Example 9

N-[2-(N'-benzylpiperidino-4) ethyl]-1,8-naphthalimide



2 g of 1,8-naphthalic anhydride and 2.2 g of 4-(2-aminoethyl) benzylpiperidine are heated and refluxed in a

- 27 -

solvent of n-butanol for 6 hours.

After allowing to stand for cooling, 50 ml of water is added to the reaction solution followed by extracting with chloroform. The resulting chloroform layer is dried  
5 over potassium carbonate, which is then concentrated to dryness under reduced pressure.


The oily residue thus obtained is purified with chloroform as a distilling solvent by the use of a silica gel column. The purified substance is formed into a hydrochloride using  
10 a 10% hydrochloric acid-ethanol solution. Upon recrystallization from ethanol-methanol, 2.0 g of the object compound is obtained (yield: 46%).

NMR ( $\delta$  value, DMSO):

2.84 (2H, d), 3.45 (2H, s), 4.4 (2H, q),  
15 7.24 (5H, s), 7.4 (2H, q), 8.1 (2H, dd),  
8.5 (2H, dd)

#### Example 10

3-[2-(1-benzyl-4-piperidino) ethyl]-2,4-(1H,3H)-  
quinazolidione . hydrochloride



A mixture of 50 g of 2-[(ethoxycarbonyl) amino] benzoic acid methyl and 49 g of 4-(2-aminoethyl)-benzylpiperidine is heated to 190 to 200 °C with stirring, and methanol and ethanol formed are distilled off.

- 5 After heating and stirring for about 6 hours, the mixture is purified using a silica gel column in a similar manner as in Example 1 and is formed into a hydrochloride.

Upon recrystallization from ethanol-water, 13.3 g of the titled compound is obtained (yield: 14.8%).

- 10 Melting point (°C): 208-210

Elemental analytical values:  $C_{22}H_{25}N_3O_2 \cdot HCl$

	C	H	N
Theoretical value (%)	66.07	6.55	10.51
Found value (%)	66.02	6.64	10.75

- 15 Examples 11 to 137

Compounds prepared using similar procedures as in Examples 1 to 10 are shown in Table 1 and Table 2.

The compounds shown in Tables 1 and 2 are represented by the formula

20





Table 1  
R'-X--R'

























Example	R'	X		R'	Melting point (°C)	Solvent for recrystallization	Molecular formula	Elemental analytical value (%)			Preparation method
								C	H	N	
11		-O(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	137~138.5	EtOH-IPe	C <sub>21</sub> H <sub>21</sub> N <sub>3</sub> O <sub>3</sub> · 1/2 H <sub>2</sub> O · HCl	62.98 62.85	6.74 6.65	7.34 7.16	A
12		-O(CH <sub>2</sub> ) <sub>2</sub> -		H	203~204	EtOH	C <sub>21</sub> H <sub>21</sub> N <sub>3</sub> O <sub>3</sub> · HCl	54.44 54.30	6.60 6.64	9.77 9.57	A
13		-O(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	100~101	EtOH-IPe	C <sub>21</sub> H <sub>21</sub> N <sub>3</sub> O <sub>3</sub>	60.73 60.50	7.34 7.61	7.29 7.30	A
14		-O(CH <sub>2</sub> ) <sub>2</sub> -			170~176	EtOH-IPe	C <sub>21</sub> H <sub>21</sub> N <sub>3</sub> O <sub>3</sub> · HCl	58.08 58.01	6.20 6.11	6.16 6.22	A
15		-O(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	179~180	EtOH	C <sub>21</sub> H <sub>21</sub> N <sub>3</sub> O <sub>3</sub> · HCl	69.44 69.51	6.69 6.72	6.06 6.11	E
16		-O(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	135~137	Acetone IPe	C <sub>21</sub> H <sub>21</sub> N <sub>3</sub> O <sub>3</sub> · HCl	68.21 68.01	5.93 6.02	5.60 5.50	E
17		-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -		-CH <sub>2</sub> - 	214~217	EtOH-H <sub>2</sub> O	C <sub>21</sub> H <sub>21</sub> N <sub>3</sub> O <sub>3</sub> · HCl · 1/2 H <sub>2</sub> O	64.12 64.12	6.53 6.50	7.12 7.24	E
18		-O(CH <sub>2</sub> ) <sub>2</sub> -			159~161	EtOH-IPe	C <sub>21</sub> H <sub>21</sub> N <sub>3</sub> O <sub>3</sub> · HCl	62.60 62.39	6.75 6.69	5.85 5.99	A

Table 1 (Continued)























Example	R'	X		R'	Melting point (°C)	Solvent for recrystallization	Molecular formula	Elemental analytical value (%)			Preparation Method
								C	II	N	
19		-COIII(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	107~113	EtOH Ether	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub> S · HCl	62.77 62.71	6.94 6.91	6.65 6.59	B
20		-COIII(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	106~110	EtOH Ether	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub> · HCl	66.26 66.15	7.01 7.10	6.72 6.61	B
21		-COIII(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	156~158	—	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub> · HCl	63.80 63.74	6.75 6.78	6.47 6.61	B
22		-COIII(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	166~168	—	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub> · HCl	78.47 76.57 (2/3 H <sub>2</sub> O) 76.54	7.53 7.58 7.62	6.54 6.24 6.30	B
23		-COIII(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	245~247 (Decomposition)	EtOH Ether	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub> S · HCl · 1/2 H <sub>2</sub> O	62.45 61.99	7.26 7.03	5.60 5.68	B
24		-COIII(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	182~184	EtOH	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub> · HCl	68.29 68.33	7.03 7.16	7.24 7.41	B
25		-COIII(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	255~257 (Decomposition)	EtOH	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub> · HCl	71.23 71.51	5.98 5.80	5.73 5.81	B

Table 1 (Continued)
























Example	R <sup>1</sup>	X			Melting point (°C)	Solvent for recrystallization	Molecular formula	Elemental analytical values			Preparation method
								Upper column: Theoretical value (%)	Bottom column: Found value (%)		
26		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	225~226	EtOH	C <sub>21</sub> H <sub>23</sub> N <sub>2</sub> O <sub>2</sub> · HCl	70.79 70.91	6.96 7.03	8.54 8.61	B
27		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	234~235	EtOH Ether	C <sub>21</sub> H <sub>21</sub> N <sub>2</sub> O · HCl	69.42 69.31	7.09 7.00	10.56 10.46	B
28		CH <sub>3</sub> -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	183~184	—	C <sub>21</sub> H <sub>22</sub> N <sub>2</sub> O · HCl	70.85 70.87	7.84 7.75	7.51 7.54	B
29		CH <sub>3</sub> -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	174~175	—	C <sub>21</sub> H <sub>23</sub> N <sub>2</sub> O <sub>2</sub> · HCl	65.93 65.98	7.46 7.44	6.69 6.65	B
30		CH <sub>3</sub> -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	164~165	—	C <sub>21</sub> H <sub>23</sub> N <sub>2</sub> O <sub>2</sub> · HCl	69.47 69.34	7.53 7.53	6.75 6.68	B
31		CH <sub>3</sub> -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	185~186	—	C <sub>21</sub> H <sub>23</sub> N <sub>2</sub> O <sub>2</sub> · HCl	70.49 70.18	7.96 7.83	6.32 6.54	B
32		CH <sub>3</sub> -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	195~196	—	C <sub>21</sub> H <sub>23</sub> N <sub>2</sub> O · HCl	74.90 74.60	7.41 7.46	6.24 6.28	B



Table I (Continued)

























Example	R <sup>1</sup>	X		R <sup>2</sup>	(°C)	Solvent for recrystallization	Molecular formula	Elemental analytical values			Preparation Method
								Upper column: Theoretical value (%)	Bottom column: Found value (%)		
33		$\text{CH}_3$ -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	235~236	EtOH	$\text{C}_{22}\text{H}_{24}\text{O}_2 \cdot \text{HCl} \cdot \frac{1}{2}\text{H}_2\text{O}$	70.37 70.12	6.30 6.23	5.47 5.28	B
34		-COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	119~123	EtOH IPG	$\text{C}_{22}\text{H}_{24}\text{O}_2 \cdot \text{HCl}$	63.91 63.88	6.97 6.90	11.10 11.16	B
35		$\text{C}_2\text{H}_5$ -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	203~204	—	$\text{C}_{23}\text{H}_{26}\text{O}_2 \cdot \text{HCl}$	70.00 70.00	7.75 7.80	6.53 6.47	B
36		-COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	161~164	—	$\text{C}_{21}\text{H}_{22}\text{O}_2$	67.98 67.91	8.56 8.48	16.99 16.76	B
37		-COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	106~110	MeOH Ether	$\text{C}_{23}\text{H}_{26}\text{O}_2 \cdot 2\text{HCl}$	63.51 63.35	6.47 6.59	10.50 10.66	B
38		-COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	127~129	MeOH IPG	$\text{C}_{23}\text{H}_{26}\text{O}_2 \cdot 3\text{HCl}$	58.09 58.18	5.92 5.99	12.09 11.01	B
39		-COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	111~122	Acetone IPG	$\text{C}_{23}\text{H}_{26}\text{O}_2 \cdot 2\text{HCl}$	60.66 60.88	6.22 6.01	7.06 7.59	B
40		$\text{CH}_3$ -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	135~136	Acetone n-Hexane	$\text{C}_{23}\text{H}_{26}\text{O}_2 \cdot \text{HCl}$	69.20 68.89	6.97 6.87	5.38 5.09	B

Table I (continued)

Example	R'	X		R'	Melting point (°C)	Solvent for recrystallization	Molecular formula	Elemental analytical values			Preparation method
								Upper column: Theoretical value(%)	Bottom column: Found value(%)	N	
41		$\text{CH}_3$ -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	176~178	—	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O <sub>4</sub> S	68.75	67.76	5.53	B
42		$\text{CH}_3$ -COH(CH <sub>2</sub> ) <sub>2</sub> -		—	149~151	EtOH PE	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O <sub>4</sub> S	65.97	7.05	6.99	B
43		$\text{CH}_3$ -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	163~164	EtOH PE	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O <sub>4</sub> S	68.75	6.76	5.53	B
44		$\text{CH}_3$ -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> CH <sub>2</sub> OH	114~116	EtOH PE	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O <sub>4</sub> · HCl	69.55	6.23	8.11	B
45		-SO <sub>2</sub> NH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	122~123	EtOH	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O <sub>4</sub> S	61.68	6.47	7.19	B
46		—		-CH <sub>2</sub> -	254~255	EtOH	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O <sub>4</sub> · HCl	59.78	5.02	10.46	C
47		—(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	177~179	EtOH Ether	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O <sub>4</sub> · HCl	61.47	5.63	9.77	C
								59.99	5.13	10.41	
								61.59	5.66	9.51	

Table 1 (continued)














Example	R'	X		R'	Melting point (°C)	Solvent for recrystallization	Molecular formula	Elemental analytical values			Preparation method
								Upper column: Theoretical value (%)	Bottom column: Found value (%)		
48		$-(CH_2)_2-$		$-CH_2-$ 	116~110	NaOH 1PE	$C_{15}H_{13}N_2O_4 \cdot HCl$	61.47 61.30	5.63 5.61	9.77 9.07	C
49		$-(CH_2)_2-$		$-CH_2-$ 	246~248 (Decomposition)	EtOH Ether	$C_{17}H_{15}N_2O_4 \cdot HCl$	69.11 68.96	6.00 6.12	8.34 8.39	C
50		$-(CH_2)_2-$		$-CH_2-$ 	230~239 (Decomposition)	NaOH Ether	$C_{17}H_{15}N_2O_4 \cdot HCl$	65.22 65.43	6.39 6.28	9.51 9.60	C
51		$-(CH_2)_2-$		$-CH_2-$ 	227~228 (Decomposition)	EtOH Ether	$C_{17}H_{15}N_2O_4 \cdot HCl$	69.55 69.29	6.23 6.34	8.11 8.08	C

Table 1 (Continued)


















Example	R'	X			R'	Melting point (°C)	Solvent for recrystallization	Molecular formula	Elemental analytical values			Preparation method
									Upper column: Theoretical value (%)	Bottom column: Found value (%)	N	
52		-(CH <sub>2</sub> ) <sub>2</sub> -			C <sub>17</sub> H <sub>13</sub> O <sub>3</sub>	161~164	H <sub>2</sub> O/Ether	C <sub>17</sub> H <sub>13</sub> O <sub>3</sub> · HCl	66.80 66.85	7.27 7.05	8.66 8.64	C
53		-(CH <sub>2</sub> ) <sub>2</sub> -			C <sub>17</sub> H <sub>13</sub> O <sub>3</sub>	162~163	—	C <sub>17</sub> H <sub>13</sub> O <sub>3</sub> · HCl	71.22 70.98	5.98 5.96	5.73 5.79	C
54		-(CH <sub>2</sub> ) <sub>2</sub> -			C <sub>17</sub> H <sub>13</sub> O <sub>3</sub>	221~222 (Decomposition)	H <sub>2</sub> O/EPE	C <sub>17</sub> H <sub>13</sub> O <sub>3</sub> · HCl	66.57 66.42	6.56 6.59	6.75 6.67	C
55		-(CH <sub>2</sub> ) <sub>2</sub> -			C <sub>17</sub> H <sub>13</sub> O <sub>3</sub>	221~224 (Decomposition)	H <sub>2</sub> O/EPE	C <sub>17</sub> H <sub>13</sub> O <sub>3</sub> · HCl	65.91 65.77	6.29 6.34	6.99 6.69	C
56		-(CH <sub>2</sub> ) <sub>2</sub> -			C <sub>17</sub> H <sub>13</sub> O <sub>3</sub>	222~223 (Decomposition)	H <sub>2</sub> O/EPE	C <sub>17</sub> H <sub>13</sub> O <sub>3</sub> · HCl	61.47 61.33	5.83 5.92	6.24 6.31	C

Table 1 (Continued)

















Example	R'	X		R'	Melting point (°C)	Solvent for recrystallization	Molecular formula	Elemental analytical values			Preparation method
								Upper column: Theoretical value (%)	Bottom column: Found value (%)	N	
57		$-(CH_2)_2-$		$-CH_2-$ 	235~236 (Decomposition)	H <sub>2</sub> O/H <sub>2</sub>	$C_{14}H_{12}H_{12}O_2 \cdot HCl$	71.80	6.26	6.44	C
58		$-(CH_2)_2-$		$-CH_2-$ 	170~181	Acetone H <sub>2</sub> O	$C_{14}H_{12}H_{12}O_2 \cdot 2HCl$	61.77 61.46	6.17 6.35	6.06 7.17	C
59		$-(CH_2)_2-$		$-CH_2-$ 	161~163	Acetone H <sub>2</sub> O	$C_{14}H_{12}H_{12}O_2 \cdot 2HCl$	61.77 61.51	6.17 6.22	6.86 7.21	C
60		$-(CH_2)_2-$		$-CH_2-$ 	210~214 (Decomposition)	H <sub>2</sub> O/H <sub>2</sub> H <sub>2</sub> O	$C_{14}H_{12}H_{12}O_2 \cdot HCl$	69.25 68.96	6.82 6.80	7.02 7.02	C
61		$-(CH_2)_2-$		$-CH_2-$ 	266~268	H <sub>2</sub> O/H <sub>2</sub> H <sub>2</sub> O	$C_{14}H_{12}H_{12}O_2 \cdot HCl$	69.47 69.31	7.53 7.46	6.72 6.63	C

Table 1 (Continued)















Example	$\eta'$	$\chi$	$\eta'$	$\eta'$	Melting point (°C)	Solvent for recrystallization	Molecular formula	Elemental analytical values			Preparation Method
								Upper column: Theoretical value (%)	Bottom column: Found value(%)		
								C	H	N	
62		$-(CH_2)_2-$		$-CH_2-$ 	198~204 (Decomposition)	$CHCl_3$ Ether	$C_{11}H_{11}O_4 \cdot HCl$	62.23 61.94	5.90 5.95	9.47 9.15	C
63		$-(CH_2)_2-$		$-CH_2-$ 	104~100 (Decomposition)	$EtOH$ Ether	$C_{11}H_{11}O_4 \cdot 2HCl$	60.55 60.21	6.24 6.11	9.63 9.57	C
64		$-(CH_2)_2-$		$-CH_2-$	256~258 (Decomposition)	$EtOH$ PE	$C_{11}H_{11}O_4 \cdot HCl$	54.31 54.29	5.70 5.71	11.00 11.81	C
65		$-(CH_2)_2-$		$-CH_2-$ 	205~206	—	$C_{12}H_{11}O_4 \cdot HCl$	60.06 60.09	5.76 5.73	9.14 9.00	C
66		$-(CH_2)_2-$		$-CH_2-$ 	220~231	—	$C_{11}H_{11}O_4 \cdot Cl \cdot HCl$	56.91 56.91	4.99 5.02	9.05 9.03	C

Table 1 (Continued)











Example	R'	X		R'	Melting point (°C)	Solvent for recrystallization	Molecular formula	Elemental analytical values Upper column: Theoretical - Bottom column: Found value (%)	Preparation Method
								C H N	
67		$-\text{CH}_2\text{CH}_2(\text{CH}_2)_2-$		$-\text{CH}_2-$ 	273~276 (Decomposition)	EtOH Ether	$\text{C}_{17}\text{H}_{17}\text{N}_2 \cdot 2\text{HCl}$	66.14 7.93 7.35 65.73 7.79 7.28	A
68		$-(\text{CH}_2)_2-$		$-\text{CH}_2-$ 	212~213	EtOH	$\text{C}_{17}\text{H}_{15}\text{O}_2\text{Cl} \cdot \text{HCl}$	60.83 5.80 9.67 60.61 5.93 9.76	D
69		$-(\text{CH}_2)_2-$		$-\text{CH}_2-$ 	221~223	EtOH Ether	$\text{C}_{17}\text{H}_{15}\text{O}_2 \cdot \text{HCl}$	66.74 6.82 10.15 66.92 6.71 10.04	D

Table 2



















Example	$R^1$	X	$R^2$	Molecular formula	$^1H$ -NMR (CDCl <sub>3</sub> , ppm)	Preparation method
70		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		C <sub>11</sub> H <sub>12</sub> N <sub>2</sub> O	1.16~2.04(9H, m), 2.7~2.96(2H, d), 3.20~3.56(4H, m), 6.0(1H, s), 7.2~7.76(10H, m)	B
71		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		C <sub>11</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub>	1.00~2.12(9H, m), 2.04~2.96(2H, d), 3.30~3.6(2H, m), 3.52(2H, s), 6.40(1H, b.t.), 7.3(5H, s), 7.9(2H, d), 8.24(2H, d)	B
72		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		C <sub>11</sub> H <sub>12</sub> N <sub>2</sub> O	1.1~2.06(9H, m), 2.0~2.9(2H, b.d), 3.32~3.56(4H, s.t.m), 6.36(1H, t), 7.26(5H, s), 7.64(2H, d), 7.84(2H, d)	B
73		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		C <sub>11</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub>	1.24~2.03(9H, m), 2.0~2.86(2H, d), 3.26~3.48(2H, m), 3.46(2H, s), 3.0(3H, s), 6.14(1H, b.t), 6.9~7.3(9H, m)	B
74		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		C <sub>11</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub>	1.23~2.04(9H, m), 2.76~2.80(2H, b.d), 3.35~3.58(4H, s.t), 3.84(9H, s), 6.38(1H, b.t), 7.0(2H, s), 7.28(5H, s)	B
75		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		C <sub>11</sub> H <sub>12</sub> N <sub>2</sub> OC	2.9(2H, d), 3.45(3H, s, d), 6.26(1H, b.t), 7.26(5H, s), 7.3(2H, d), 7.7(2H, d)	B
76		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		C <sub>11</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub>	2.9(2H, d), 3.46(3H, s, d), 3.78(6H, s), 6.12(1H, b.t), 6.5(1H, t), 6.82(2H, d), 7.26(5H, s)	B
77		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		C <sub>11</sub> H <sub>12</sub> N <sub>2</sub> O	2.7(2H, d), 2.8(6H, s), 3.34(2H, s), 6.46(2H, d), 6.5(1H), 7.12(5H, s), 7.59(2H, d)	B



Table 2 (Continued)
























Example	R'	X		R''	Molecular formula	<sup>1</sup> H-NMR (CDCl <sub>3</sub> , ppm)	Preparation method
70		-COOH (CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>14</sub> O <sub>3</sub>	1.1 ~ 2.0 (9H, m), 2.76 ~ 2.86 (2H, bd), 3.26 ~ 3.40 (4H, m), 5.94 (2H, s), 6.32 (1H, bt), 6.72 (1H, d), 7.24 (2H, s)	B
79	CH <sub>2</sub> S- 	-COOH (CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>12</sub> O <sub>3</sub> S	1.02 ~ 2.04 (9H, m), 2.8 ~ 2.9 (2H, bd), 2.8 ~ 2.9 (2H, bd), 3.2 ~ 3.5 (4H, s, m), 6.26 (1H, bt), 7.10 (2H, d), 7.20 (5H, s), 7.64 (2H, d)	B
80	CH <sub>2</sub> SO <sub>2</sub> - 	-COOH (CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>12</sub> O <sub>3</sub> S	1.12 ~ 2.06 (9H, m), 2.8 ~ 2.9 (2H, bd), 3.04 (2H, s), 3.36 ~ 3.50 (2H, m), 3.48 (2H, s), 6.3 (1H, bt), 7.26 (5H, s), 7.9 (4H, s)	B
81	CH <sub>2</sub> - 	-COOH (CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>12</sub> O <sub>3</sub>	1.3 ~ 2.08 (9H, m), 2.86 (2H, bd), 3.46 (2H, s), 6.0 (1H, bs), 7.16 (5H, s), 7.64 (2H, d), 7.82 (2H, d)	B
82	CH <sub>2</sub> O- 	-COOH (CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>12</sub> O <sub>3</sub>	1.2 ~ 2.20 (9H, m), 2.98 (2H, bd), 3.42 (3H, m), 3.60 (2H, s), 7.30 (5H, s), 7.92 (4H, s), 10.02 (1H, s)	B
83	HO- 	-COOH (CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>12</sub> O <sub>3</sub>	1.2 ~ 2.04 (9H, m), 2.84 (2H, bd), 3.30 (2H, m), 3.46 (2H, s), 6.52 (1H, bs), 6.66 (2H, d), 7.22 (5H, s), 7.52 (2H, d), 8.30 (1H, s)	B
84	 -CH <sub>2</sub> O- 	-COOH (CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>12</sub> O <sub>3</sub>	1.1 ~ 2.04 (9H, m), 2.84 (2H, bd), 3.44 (4H, s, m), 5.06 (2H, m), 6.00 (1H, bt), 6.92 (2H, d), 7.14 ~ 7.44 (10H, m), 7.68 (2H, d)	B

Table 2 (Continued)

Example	R <sup>1</sup>	X	A	R <sup>2</sup>	Molecular formula	<sup>1</sup> H-NMR (CDCl <sub>3</sub> , ppm)	Preparation method
85		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	C <sub>14</sub> H <sub>18</sub> O <sub>2</sub>	1.06~2.04 (15H, m), 2.0 (2H, bd), 3.10~3.60 (1H, stw), 6.96 (1H, bt), 7.22 (1H, std), 7.66 (2H, d)	B
86		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	C <sub>17</sub> H <sub>18</sub> O <sub>2</sub>	1.1~2.1 (15H, m), 2.76~2.88 (2H, bd), 3.3~3.5 (2H, m), 3.44 (2H, s), 4.8 (1H, m), 5.7~6.04 (2H, m), 6.88 (2H, d), 7.28 (5H, m), 7.64 (2H, d)	B
87		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	C <sub>15</sub> H <sub>18</sub> O <sub>2</sub>	1.3~2.0 (7H, m), 2.7~3.4 (6H, m), 4.14~4.36 (5H, m), 7.3~8.04 (9H, m), 8.40 (1H, m), 10.00 (1H, m)	B
88		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	C <sub>18</sub> H <sub>18</sub> O <sub>2</sub> S	1.0~1.98 (9H, m), 2.6~2.86 (2H, m), 3.00~3.5 (1H, st), 4.20 (2H, s), 7.2~7.4 (12H, m), 7.7 (2H, d), 8.32 (1H, bt)	B
89		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	C <sub>17</sub> H <sub>18</sub> O <sub>2</sub> S	1.2~2.3 (17H, m), 3.00~3.2 (2H, bd), 2.5~2.7 (5H, m), 3.1 (1H, m), 7.06~7.9 (9H, m)	B
90		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	C <sub>17</sub> H <sub>18</sub> O <sub>2</sub>	1.24~2.04 (9H, m), 2.76~2.88 (2H, bd), 3.3~3.5 (6H, stw), 6.1 (1H, bt), 6.88~7.74 (14H, m)	B
91		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	C <sub>18</sub> H <sub>18</sub> O <sub>2</sub> S	1.22~2.06 (9H, m), 2.8~2.92 (2H, bd), 3.30 (3H, s), 3.44 (2H, s), 3.20~3.44 (2H, m), 4.94 (2H, s), 6.36 (1H, bt), 7.26 (5H, s), 7.4 (2H, d), 7.64 (2H, d)	B
92		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -	C <sub>17</sub> H <sub>18</sub> O <sub>2</sub>	1.24~2.06 (9H, m), 2.88 (2H, bd), 3.40 (2H, s), 3.48 (2H, s), 5.11 (2H, s), 5.94 (1H, bs), 6.94 (2H, d), 7.20 (5H, s), 7.32 (2H, d), 7.7 (2H, d), 8.6 (2H, d)	B

Table 2 (Continued)





























Example	R <sup>1</sup>	X		R <sup>2</sup>	Molecular formula	<sup>1</sup> H-NMR (CDCl <sub>3</sub> , ppm)	Preparation method
93		-COH(CCl <sub>4</sub> ), -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>12</sub> O	2.9 (2H, d), 3.40 (3H, s, d), 6.7 (1H, h), 7.20 (6H, s), 7.1 (1H, s), 7.65 (1H, s), 7.9 (1H, s)	B
94		-COH(CCl <sub>4</sub> ), -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>11</sub> Cl	1.26 ~ 2.05 (9H, m), 2.0 ~ 2.9 (2H, h), 3.32 ~ 3.54 (4H, s, m), 6.56 (1H, h), 7.20 (5H, s), 7.34 (1H, d), 8.02 (1H, d), 8.7 (1H, d)	B
95		-COH(CCl <sub>4</sub> ), -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>11</sub> O	1.3 ~ 2.07 (9H, m), 2.0 ~ 2.92 (2H, h), 2.30 ~ 2.6 (4H, s, m), 7.27 (5H, s), 7.7 (1H, h), 8.5 (1H, e), 8.7 (1H, d), 9.30 (1H, d)	B
96		-COH(CCl <sub>4</sub> ), -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>11</sub> O	2.16 (2H, d), 3.40 (4H, s, d), 6.44 (1H, h), 7.26 (5H, s), 7.4 ~ 7.8 (3H), 8.4 (2H), 8.74 (1H, d)	B
97		-COH(CCl <sub>4</sub> ), -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>11</sub> O	2.24 ~ 2.56 (9H, m), 3.56 (2H, s), 4.0 (2H, t), 5.20 (1H, t), 6.64 (1H, bs), 7.31 (5H, s), 7.6 (2H, dd), 8.66 (2H, dd)	B
98		 -COH(CCl <sub>4</sub> ), -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>11</sub> O <sub>2</sub> · HCl	0.8 ~ 2.10 (11H, m), 2.49 ~ 2.88 (7H, m), 7.04 ~ 7.44 (14H, m)	B
99	 Cl <sub>2</sub> SO <sub>2</sub> -	 -COH(CCl <sub>4</sub> ), -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>11</sub> O <sub>2</sub> S · HCl	0.92 ~ 2.10 (9H, m), 2.44 ~ 3.6 (7H, m), 3.0 (3H, s), 3.44 (2H, h), 7.14 (5H, s), 7.52 (2H, d), 7.96 (2H, d)	B
100		 -COH(CCl <sub>4</sub> ), -		-CH <sub>2</sub> - 	C <sub>12</sub> H <sub>11</sub> O	1.02 ~ 2.10 (12H, m), 2.7 ~ 3.7 (9H, m), 0.2 ~ 0.20 (7H, d, s), 0.60 (2H, d)	B

Table 2 (Continued)























Example	R'	X	(A)	R'	Molecular formula	<sup>1</sup> H-NMR (CDCl <sub>3</sub> , ppm)	Preparation method
101		 - -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>17</sub> H <sub>21</sub> N <sub>2</sub> O	1.04~2.04 (9H, m), 2.7~3.7 (6H, m), 4.4 (2H, s), 4.74 (2H, s), 7.00~7.6 (12H, m), 0.64 (2H, bs)	B
102		-COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>17</sub> H <sub>21</sub> N <sub>2</sub> O	1.14~2.04 (11H, m), 2.70~2.9 (2H, bs), 3.16~3.76 (4H, s, d), 6.5 (1H, s), 7.26 (5H, s), 7.56 (2H, d), 0.60 (2H, d)	B
103		-COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>17</sub> H <sub>21</sub> N <sub>2</sub> O	1.3~2.1 (9H, m), 2.76~2.88 (2H, m), 3.16~3.58 (4H, m, s), 6.84 (1H, s), 7.24 (2H, d), 7.6 (2H, d), 0.48 (2H, d), 0.60 (2H, d)	B
104		-COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>17</sub> H <sub>21</sub> N <sub>2</sub> O <sub>2</sub> Cl <sub>2</sub>	2.0 (2H, d), 3.4 (4H, s, t), 7.5~7.1 (3H), 7.62 (2H), 0.64 (2H)	B
105		-COH(CH <sub>2</sub> ) <sub>2</sub> -			C <sub>17</sub> H <sub>21</sub> N <sub>2</sub> O	1.22~1.96 (7H, m), 2.7~3.0 (2H, m), 3.4~3.62 (2H, m), 3.0~3.65 (2H, m), 6.62 (3H, m, s), 7.50 (2H, d), 0.7 (2H, d)	B
106		-COH(CH <sub>2</sub> ) <sub>2</sub> -			C <sub>17</sub> H <sub>21</sub> N <sub>2</sub> O <sub>2</sub>	1.04~2.16 (9H, m), 2.82 (2H, d), 3.32~3.62 (2H, m), 3.46 (2H, s), 6.32~6.58 (1H, m), 7.16~8.56 (13H, m)	B
107		-CH <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>17</sub> H <sub>21</sub> N <sub>2</sub> O <sub>2</sub>	1.26~2.06 (9H, m), 2.70~2.9 (2H, bs), 3.45 (2H, s), 3.62 (2H, d), 7.24 (5H, s), 7.95~8.04 (1H, m), 8.5~8.6 (2H, m)	C

Table 2 (Continued)
















Example	R <sup>1</sup>	X	(A)	R <sup>2</sup>	Molecular formula	<sup>1</sup> H-NMR (CDCl <sub>3</sub> , ppm)	Preparation method
108		-(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub>	1.26~2.04 (9H, m), 2.04 (2H, bd), 3.44 (2H, s), 3.60 (2H, t), 7.24 (5H, s), 7.7 (4H, m)	C
109		-(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub>	1.20~2.04 (9H, m), 2.05 (2H, bd), 3.46 (2H, s), 3.62 (2H, t), 5.17 (2H, s), 6.70 (1H, d), 7.1 (1H, d), 7.25 (5H, s), 7.36 (1H, t)	C
110		-(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub>	1.26~2.04 (9H, m), 2.04 (2H, bd), 3.48 (2H, s), 3.6 (2H, t), 5.31 (2H, s), 6.76 (1H, dd), 6.98 (1H, d), 7.27 (5H, s), 7.54 (1H, d)	C
111		-(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub>	1.20~2.06 (9H, m), 2.06 (2H, bd), 3.50 (2H, s), 3.60 (2H, t), 7.28 (5H, s), 7.50 (3H, m), 7.66 (3H, m), 8.1 (2H, m), 8.36 (1H, s)	C
112		-(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>11</sub> N <sub>2</sub> O <sub>2</sub>	2.04 (2H, d), 2.44 (2H, s), 2.7 (2H, t), 7.24 (10H, s), 8.2 (2H, s)	C

Table 2 (Continued)






















Example	R <sup>1</sup>	X	(A)	R <sup>2</sup>	Molecular formula	<sup>1</sup> H-NMR (CDCl <sub>3</sub> , ppm)	Preparation method
113		-(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>13</sub> H <sub>17</sub> O	1.0 ~ 2.2 (8H, m), 2.41 (3H, s), 2.7 ~ 3.1 (2H, m), 3.5 (2H, s), 4.11 (2H, t), 7.8 ~ 8.9 (8H, m)	D
114		-(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>13</sub> H <sub>17</sub> O	0.9 ~ 2.2 (8H, m), 2.7 ~ 3.0 (2H, m), 3.48 (2H, s), 4.06 (2H, t), 7.9 ~ 8.7 (8H, m), 8.0 ~ 8.2 (1H, m)	D
115		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>17</sub> H <sub>19</sub> ClNOS 1/2 H <sub>2</sub> O	0.92 ~ 2.04 (9H, m), 2.64 ~ 3.00 (2H, bd), 3.12 ~ 3.60 (2H, m), 3.40 (2H, s), 4.96 (2H, s), 6.48 ~ 6.76 (1H, bt), 7.08 ~ 7.84 (9H, m)	B
116		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>18</sub> H <sub>19</sub> ClOS 3/2 H <sub>2</sub> O	1.0 ~ 2.10 (9H, m), 2.70 ~ 2.96 (2H, bd), 3.20 ~ 3.60 (2H, m), 3.46 (2H, s), 4.00 (2H, s), 6.20 (1H, bs), 6.84 ~ 7.80 (14H, m)	B
117		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>17</sub> H <sub>19</sub> Cl	1.20 (3H, s), 1.20 (3H, s), 1.00 ~ 2.00 (9H, m), 2.64 ~ 3.02 (3H, m), 3.45 (2H, s), 3.24 ~ 3.56 (2H, m), 6.36 (1H, bt), 7.12 ~ 7.76 (9H, m)	B
118		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>17</sub> H <sub>19</sub> Cl	1.00 ~ 3.00 (14H, m), 2.64 ~ 2.96 (2H, bd), 3.44 (2H, s), 3.32 ~ 3.48 (2H, m), 3.80 (3H, s), 6.76 ~ 7.52 (9H, m)	D
119		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>17</sub> H <sub>19</sub> Cl	1.00 ~ 3.08 (14H, m), 2.60 (3H, s), 3.24 ~ 3.62 (2H, bd), 3.46 (2H, s), 6.16 (1H, bt), 7.26 (5H, s), 7.08 ~ 8.00 (14H, bd)	B

Table 2 (Continued)


























Example	R'	X		R''	Molecular formula	<sup>1</sup> H-NMR (CDCl <sub>3</sub> , ppm)	Preparation method
120		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O · HCl	1.04 ~ 2.12 (14H, m), 2.36 (3H, s), 2.60 ~ 2.96 (2H, bd), 3.20 ~ 3.56 (2H, m), 3.44 (2H, s), 6.00 (1H, bt), 7.00 ~ 7.70 (9H, m)	B
121		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>13</sub> H <sub>15</sub> N <sub>2</sub> O <sub>2</sub> · HCl	1.10 ~ 2.00 (14H, m), 2.60 ~ 2.90 (2H, bd), 3.20 ~ 3.56 (2H, m), 3.46 (2H, s), 3.80 (3H, s), 3.89 (3H, s), 5.94 (1H, bt), 6.72 ~ 8.44 (8H, m)	B
122		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O <sub>2</sub> · HCl	0.72 ~ 2.40 (9H, m), 2.40 ~ 3.00 (2H, bd), 3.40 (2H, s), 3.20 ~ 3.60 (2H, m), 6.40 ~ 6.60 (1H, bt), 7.00 ~ 7.92 (14H, m)	B
123		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O · HCl	1.12 ~ 3.12 (9H, m), 2.60 ~ 3.00 (2H, bd), 3.32 ~ 3.62 (2H, m), 3.46 (2H, s), 6.06 (1H, bt), 7.12 ~ 7.00 (14H, m)	B
124		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O <sub>2</sub> · HCl	0.92 ~ 3.00 (14H, m), 2.64 ~ 2.90 (2H, bd), 3.45 (2H, s), 3.24 ~ 3.60 (2H, m), 6.24 (1H, bt), 7.04 ~ 8.20 (14H, m)	B
125		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O · HCl	0.96 ~ 2.00 (14H, m), 2.60 ~ 2.96 (2H, bd), 3.44 (2H, s), 3.20 ~ 3.62 (2H, m), 3.26 (1H, bt), 7.00 ~ 8.20 (14H, m)	D
126		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O · HCl	1.2 ~ 2.06 (7H, m), 2.60 ~ 2.92 (2H, bd), 3.33 ~ 3.6 (2H, s), 3.46 (2H, s), 7.27 (5H, s), 7.3 ~ 8.54 (5H, m)	B
127		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> - 	C <sub>11</sub> H <sub>13</sub> N <sub>2</sub> O <sub>2</sub> · HCl	1.04 ~ 2.10 (14H, m), 2.60 ~ 2.90 (2H, bd), 3.46 (2H, s), 3.24 ~ 3.56 (2H, m), 6.06 (1H, bt), 6.00 ~ 7.04 (9H, m)	D

Table 2 (Continued)

































Example	R'	X			R <sup>2</sup>	Molecular formula	<sup>1</sup> H-NMR (CDCl <sub>3</sub> , ppm)	Preparation method
128		$\text{CH}_3$   -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -		C <sub>17</sub> H <sub>15</sub> N, O, 0.1 · HCl	0.96~3.08 (14H, m), 2.60~3.03 (2H, bd), 2.97 (3H, s), 3.03~3.60 (2H, m), 3.45 (3H, s), 3.79 (3H, s), 6.72~7.40 (9H, m)	B
129		$\text{CH}_3$   -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -		C <sub>17</sub> H <sub>15</sub> N, O, 0.1 · HCl · 1/2 H <sub>2</sub> O	1.04~2.04 (9H, m), 2.80 (2H, broad d), 2.90 (3H, d), 3.20~3.48 (2H, m), 3.24 (2H, s), 5.06 (2H, s), 6.90 (2H, d), 7.16~7.44 (12H, m)	B
130		$\text{C}_2\text{H}_5$   -COH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -		C <sub>21</sub> H <sub>19</sub> N, O, 0.5 · HCl · 1/2 H <sub>2</sub> O	0.92~1.96 (12H, m), 2.60~3.54 (8H, m), 4.36 (2H, s), 6.96~7.44 (12H, m), 7.60 (2H, d)	B
131		-CONHCH <sub>2</sub> -		-CH <sub>2</sub> -		C <sub>17</sub> H <sub>15</sub> N, O	1.2~2.08 (7H, m), 2.82~2.95 (2H, bd), 3.27~3.4 (2H, t), 3.48 (2H, s), 5.4 (1H), 7.26 (5H, s), 7.53~7.60 (2H), 8.65~8.73 (2H)	B
132		-CONH(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -		C <sub>21</sub> H <sub>19</sub> N, O, 0.1 · 2HCl	1.04~2.20 (14H, m), 2.64~2.95 (2H, bd), 3.32~3.64 (2H, m), 3.46 (2H, s), 6.48 (1H, bt), 7.12~8.56 (13H, m)	B
133		-(CH <sub>2</sub> ) <sub>2</sub> -		-CH <sub>2</sub> -		C <sub>21</sub> H <sub>19</sub> N, O, 0.1 Cl	1.07~2.50 (9H, m), 2.71~2.82 (2H, bd), 3.36 (2H, s), 3.5~3.61 (2H, t), 7.16 (5H, s), 7.46~7.76 (3H, m)	C



Table 2 (Continued)

Example	$R^1$	X	(A)	$R^2$	Molecular formula	$^1H$ -NMR(CDCl <sub>3</sub> , ppm)	Preparation Method
134		$\begin{array}{c} \text{CON}-(\text{CH}_2)_2- \\   \\ \text{CH}_3 \end{array}$		$-\text{CH}_2-$ 	C <sub>22</sub> H <sub>27</sub> N <sub>3</sub> O <sub>3</sub>	1.05~2.05(9H, m), 2.80~3.1(5H, m), 3.50(2H, d), 7.35(5H, s), 7.60(2H, d), 8.30(2H, d),	
135		$\begin{array}{c} \text{CON}-(\text{CH}_2)_2- \\   \\ \text{CH}_2\text{CH}_3 \end{array}$		$-\text{CH}_2-$ 	C <sub>23</sub> H <sub>29</sub> N <sub>3</sub> O <sub>3</sub>	1.05~2.0(9H, m), 3.2(2H, d), 3.45(2H, d), 7.30(5H, s), 7.50(2H, d),	B
136		$\begin{array}{c} \text{CON}-(\text{CH}_2)_2- \\   \\ \text{CH}_3 \end{array}$		$-\text{CH}_2-$ 	C <sub>22</sub> H <sub>27</sub> N <sub>3</sub> O	1.0~2.0(9H, m), 2.65~3.05, (5H, m), 3.45(2H, s), 7.30(5H, s), 7.45(2H, d), 7.70(2H, d), 8.30(2H, d),	B
137		$\begin{array}{c} \text{CON}-(\text{CH}_2)_2- \\   \\ \text{CH}_2\text{CH}_3 \end{array}$		$-\text{CH}_2-$ 	C <sub>23</sub> H <sub>29</sub> N <sub>3</sub> O	1.0~2.0(9H, m), 2.8(2H, d), 3.15(2H, d), 3.45(2H, s), 7.30(5H, s), 7.35(2H, d), 7.60(2H, d),	B

In order to show the effects of the compounds of the invention in detail, part of examples of pharmacological experiments using animals will be described hereinunder.

In addition to the working examples disclosed  
5 above in view of the pharmacological tests, the compound of the invention was examined in view of effects on scopolamine-induced impairment of passive avoidance, using ddY male mice. As a result it was found that it had an excellent activity in this respect.

10

Experimental Example 1

Acetylcholinesterase Inhibitory Action using Mouse Brain Homogenate

Using a mouse brain homogenate as a source of  
15 acetylcholinesterase, the esterase activity was measured by a thiocholine method. Acetylcholine as a substrate, the compound of the invention as a substance to test and 5,5'-dithio-bis(2-nitrobenzoic acid), called also DTNB, were added to the mouse brain homogenate. After  
20 incubation, thiocholine produced was reacted with DTNB to form a yellow product. The acetylcholinesterase activity was determined by the measurement of the absorbance change of the reaction product at 412 nm.

The acetylcholinesterase inhibitory activity of  
25 the compound tested was expressed in terms of the 50% inhibitory concentration, IC50.

Results are shown in Table 3.

Table 3

0229391

Compound (Example No.)	AchE inhibitory activity IC <sub>50</sub> (μM)	Compound (Example No.)	AchE inhibitory activity IC <sub>50</sub> (μM)
4	0.0006	56	0.0033
5	0.088	57	0.0012
6	0.021	58	0.013
7	0.014	62	0.0034
8	0.08	68	0.0045
9	0.0055	71	0.055
10	0.0042	88	0.049
16	0.059	98	0.0083
21	0.021	110	0.0088
25	0.0045	111	0.00088
26	0.009	112	0.00015
32	0.0043	113	0.0047
41	0.0008	115	0.0067
47	0.009	116	0.003
48	0.0039	122	0.02
50	0.0028	123	0.0105
51	0.0022	130	0.0003
52	0.0015	133	0.0079
53	0.0024	Physo *1	0.89
54	0.008	THA *2	0.084

(Notes) #1; physostigmine

#2; 1, 2, 3, 4-Tetrahydro-9-Aminoacridine

- 51 -

## Experimental Example 2

Acute Toxicity Test for ddY Male Mouse

Acute toxicity test was carried out using ddY male mice.

5 The results are shown in Table 4.


Table 4

Compound (Example No.)	Oral to mouse (mg/kg)	
	100 mg	300 mg
4	0/3 *1	2/4 *1
5	1/4	4/4
7	-	0/3
10	3/4	-

(Notes) \*1; Denominator shows the number of Animals used, and numerator shows the number of deaths.

CLAIMS

1. A piperidine derivative having the formula  
(I) or a pharmacologically acceptable salt thereof:



5 wherein  $R^1$  denotes a univalent group derived from one  
selected among substituted or unsubstituted benzene, pyridine,  
pyrazine, indole, anthraquinone, quinoline, substituted or  
unsubstituted phthalimide, homophthalimide, pyridinecarboxylic  
acid imide, pyridine N-oxide, pyrazinedicarboxylic acid imide,  
10 naphthalenedicarboxylic acid imide, substituted or  
unsubstituted quinazolinedione, 1,8-naphthalimide, bicyclo  
[2.2.2] oct-5-ene-2,3-dicarboxylic acid imide and  
pyromerylimide,

X denotes a group of the formula  $-(CH_2)_n-$ , a group of  
15 the formula  $-O(CH_2)_n-$ , a group of the formula  $-S(CH_2)_n-$ ,  
a group of the formula  $-NH(CH_2)_n-$ , a group of the formula  
 $-SO_2NH(CH_2)_n-$ , a group of the formula  $-NH-\overset{\overset{O}{||}}{C}-(CH_2)_n-$ , a group

of the formula  $\text{-NH(CH}_2\text{)}_n\text{-C-}$ , a group of the formula


$\text{-C-O(CH}_2\text{)}_n\text{-}$ , a group of the formula  $\text{-CH}_2\text{NH(CH}_2\text{)}_n\text{-}$ , a group


5 of the formula  $\text{-C-N-(CH}_2\text{)}_n\text{-}$  (in all the above formulas,



$n$  is an integer of 1 through 7 and  $\text{R}^3$  represents a lower alkyl group or a benzyl group), a group of the formula

10  $\text{-O-CH}_2\text{CH}_2\text{CH(CH}_3\text{)-}$ , a group of the formula  $\text{-O-CH(CH}_3\text{)CH}_2\text{CH}_2\text{-}$ , a group of the formula  $\text{-O-CH}_2\text{CH}_2\text{CH=}$  or a group of the formula


$\text{-O-CH}_2\text{-CH(OH)-CH}_2\text{-}$ ,

15 the ring A denotes a group of the formula ,

a group of the formula , a group of the formula

 or a group of the formula , and

20  $\text{R}^2$  denotes a hydrogen atom, a lower alkyl group, a substituted or unsubstituted benzyl group, a substituted or

unsubstituted benzoyl group, a pyridyl group, a 2-hydroxyethyl group, a pyridylmethyl group or a group of the formula  $Z$   (wherein Z represents a halogen atom).

5        2. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which R1 is a pyridyl group.

3. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which R1  
10 is a univalent group derived from a substituted or unsubstituted benzene.

4. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which R1 is a univalent group derived from a substituted or  
15 unsubstituted phthalimide.

5. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which R1 is a univalent group derived from homophthalimide.


6. A piperidine derivative or a pharmacologically  
20 acceptable salt as claimed in Claim 1, in which R1

is a univalent group derived from a substituted or unsubstituted quinazolinedione.

7. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which X  
5 is a group of the formula:  $-(CH_2)_n-$ ,  
n being an integer of 1 to 7.


8. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which X  
is a group of the formula:

10



wherein n is an integer of 1 through 7 and  $R^3$  represents a lower alkyl group or a benzyl group.


15 9. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which the ring A is a group of the formula:




20 10. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which X  
is a group of the formula:




- 56 -




wherein n is an integer of 1 through 7 and R<sup>3</sup> represents a lower alkyl group or a benzyl group, and the ring A is a group of the formula



11. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which X is a group of the formula:




wherein n is an integer of 1 through 7 and R<sup>3</sup> represents a lower alkyl group or a benzyl group, the ring A is a group of the formula




and R<sup>2</sup> is a substituted or unsubstituted benzyl group.

12. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which R<sup>1</sup> is a univalent group derived from a substituted or unsubstituted benzene, X is a group of the formula:



wherein n is an integer of 1 through 7 and R<sup>3</sup> represents a lower alkyl group or a benzyl group, the ring A is a group of the formula




5


and R<sup>2</sup> is a substituted or unsubstituted benzyl group.

13. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which R<sup>1</sup> is a group of the formula:


10



X is a group of the formula




15 wherein R<sup>3</sup> represents a lower alkyl group, the ring A is a group of the formula




and R<sup>2</sup> is a benzyl group.

20


14. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which R<sup>1</sup> is a group of the formula:



X is a group of the formula




the ring A is a group of the formula




and R<sup>2</sup> is a benzyl group.

- 10            15. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which R<sub>1</sub> is a univalent group derived from a substituted or unsubstituted phthalimide, X is a group of the formula:



- 15    wherein n is an integer of 1 through 7, the ring A is a group of the formula




and R<sup>2</sup> is a substituted or unsubstituted benzyl group.

- 20            16. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which n is 2.

17. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which


the compound is N-[2-(N'-benzylpiperidino-4) ethyl]-4-nitrophthalimide.

18. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which R<sub>1</sub> is a univalent group derived from homophthalimide, X is a group of the formula:



wherein n is an integer of 1 through 7, the ring A is a group of the formula

10




and R<sup>2</sup> is a substituted or unsubstituted benzyl group.

19. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 18, in which n is 2.

20. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which the compound is N-[2-(N'-benzylpiperidino-4) ethyl]-homophthalimide.


21. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which R<sub>1</sub> is a univalent group derived from a substituted or unsubstituted quinazolinedione, X is a group of the formula:

25



- 60 -

wherein n is an integer of 1 through 7, the ring A is a group of the formula



5 and R<sup>2</sup> is a substituted or unsubstituted benzyl group.

22. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 21, in which n is 2.

23. A piperidine derivative or a pharmacologically acceptable salt as claimed in Claim 1, in which the  
10 compound is 3-[2-(1-benzylpiperidino) ethyl-2,4-(1H,3H)-quinazolinedione.

24. A pharmaceutical composition which comprises the piperidine derivative as defined in Claim 1 and  
15 a pharmacologically acceptable carrier.

25. The use of a compound according to claim 1 for the making of a pharmaceutical composition preventing dementias and sequelae of cerebrovascular diseases.



European Patent  
Office

# EUROPEAN SEARCH REPORT

0229391

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	EP - A2 - 0 112 776 (RHONE-POULENC) * Claims 1,5; examples 47,49, 50; abstract *	1,10, 11,16, 24	C 07 D 211/26 C 07 D 401/12 C 07 D 401/06 C 07 D 211/22 C 07 D 211/14 C 07 D 211/32 C 07 D 211/70 C 07 D 211/94 C 07 D 401/04 C 07 D 401/14 C 07 D 405/12 C 07 D 471/04 C 07 D 487/04 A 61 K 31/445 A 61 K 31/47 A 61 K 31/505
X	GB - A - 1 507 462 (ANTONIO GALLARDO) * Claims 1,21; page 1, lines 8-16; table I, compound no. 10 *	1,24	
X	GB - A - 1 268 909 (ROBINS COMPANY) * Claims 1,40; page 2, lines 4-9 *	1,3, 7,16, 24	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 07 D 211/00 C 07 D 401/00 C 07 D 405/00 C 07 D 471/00 C 07 D 487/00
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 03-04-1987	Examiner HOCHHAUSER
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	